

Figure 1. Schematic of conditional simulation process.




Figure 2. Illustration 1: Comparison of simulated and interpolated bathymetric elevations for a pair of equally likely realizations in panel (A). Panel (B) is an expanded view of the same pair as panel (A) and panel (C) is an alternative but equally likely pair form another random draw. Blocks 1 and 2 represent intervals over which mean elevation change is characterized.




Figure 3. Illustration 1: Five equally likely simulated bathymetry surfaces (Panel A) and the ensemble of 1000 surfaces defining an envelope of uncertainty about the interpolated line (Panel B).




Figure 4. Illustration 1: Histogram of elevation change for blocks 1 and 2 and blocks 1 and 2 combined. The vertical line represents the 6 inch threshold used to define elevation changes distinguishable potential offsets due to measurement bias. Negative values represent erosion and positive values represent deposition.



Figure 5. Illustration 1: Cumulative probability distribution for average elevation change fro blocks 1, 2 and block 1 and 2 combined. Probability of some erosion (i.e. negative change) is approximately 70%, 97% and 94% respectively for blocks 1, 2 and (1 and 2) combined. Yet the probability of more than 6 inches of erosion is less than 50% so these blocks would be classified as not erosional.





Figure 6. Illustration 2: Comparison of simulated and interpolated bathymetric elevations for a pair of equally likely realizations in panel (A). Panel (B) is an expanded view of the same pair as panel (A) and panel (C) is an alternative but equally likely pair form another random draw. Blocks 1 and 2 represent intervals over which mean elevation change is characterized.






Figure 7. Illustration 2: Five equally likely simulated bathymetry surfaces (Panel A) and the ensemble of 1000 surfaces defining an envelope of uncertainty about the interpolated line (Panel B).




Figure 8. Illustration 2: Histogram of elevation change for blocks 1 and 2 and blocks 1 and 2 combined. The vertical line represents the 6 inch threshold used to define elevation changes distinguishable potential offsets due to measurement bias. Negative values represent erosion and positive values represent deposition.




Figure 9. Illustration 2: Cumulative probability distribution for average elevation change fro blocks 1, 2 and block 1 and 2 combined. Probability of some erosion (i.e. negative change) is approximately 60% to 80% respectively for blocks 1, 2 and (1 and 2) combined. Yet the probability of more than 6 inches of erosion is less than 50% so these blocks would be classified as not erosional.




Figure 10. Schematic semi-variogram with definitions parameters.

Figure 11: Long flow and cross flow coordinate axes resulting from the Schwartz-Christoffel conformal mapping in a selected part of the lower Passaic River.




Figure 12a. Histogram and Summary Statistics for 1989 Bathymetry Survey




Figure 12b. Histogram and Summary Statistics for 1995 Bathymetry Survey




Figure 12c. Histogram and Summary Statistics for 1996 Bathymetry Survey




Figure 12d. Histogram and Summary Statistics for 1997 Bathymetry Survey




Figure 12e. Histogram and Summary Statistics for 1999 Bathymetry Survey




Figure 12f. Histogram and Summary Statistics for 2001 Bathymetry Survey




Figure 12g. Histogram and Summary Statistics for 2002 Bathymetry Survey




Figure 12h. Histogram and Summary Statistics for 2004 Bathymetry Survey




Figure 12i: Polynomial Regression Result for 1989 bathymetry

Multiple Regression Analysis for 1989 Bathymetry Survey Depth and Transformed River Coordinates (Note: U = Cross Flow Coordinate; V = Along Flow Coordinate)

Dependent variable: Depth

Parameter	Estimate	Standard Error	T Statistic	P-Value
CONCEAND	14 60000	0.026000	C1 700	
CONSTANT U	-14.62280 2.47516	0.236888 0.215729	-61.729 11.474	0.0000
٧	0.03028	0.002888	10.484	0.0000
U*U	0.90473	0.112363	8.052	0.0000
V*V	-0.00015	0.000008	-17.603	0.0000
υ*Δ	-0.01657	0.000966	-17.152	0.0000

Analysis of Variance

Source Su	m of Squares	Df	Mean Square	F-Ratio	P-Value
Model Residual	95633.4 611305.0	5 16611	19126.7 36.8012	519.73	0.0000

Total (Corr.) 706938.0 16616

R-squared = 13.5278 percent

R-squared (adjusted for d.f.) = 13.5018 percent

Standard Error of Est. = 6.0664

Mean absolute error = 4.79136

Durbin-Watson statistic = 0.130868 (P=0.0000)

Figure 12j: Polynomial Regression Result for 1995 bathymetry. Inset Figure is Semivariogram of Residuals

Multiple Regression Analysis for 1995 Bathymetry Survey Depth and Transformed River Coordinates (Note: U = Cross Flow Coordinate; V = Along Flow Coordinate)

Dependent variable: Depth

Parameter	Estimate	Standard Error	T Statistic	P-Value
CONSTANT	-6.01978	0.474852	-12.6772	0.0000
υ	-1.09495	0.228131	-4.7997	0.0000
V	-0.08985	0.005290	-16.9857	0.0000
U*U	5.32106	0.091953	57.8671	0.0000
Δ*Δ	0.00022	0.000014	15.9645	0.0000
$\Pi \star \Delta$	-0.00474	0.001005	-4.7121	0.0000

Analysis of Variance

Source	Sum of Squares	\mathbf{Df}	Mean Square	F-Ratio	P-Value
Model	181699.0	5	36339.8	872.21	0.0000
Residual	868240.0	20839	41.6642		

Total (Corr.) 1.04994E6 20844

R-squared = 17.3057 percent

R-squared (adjusted for d.f.) = 17.2858 percent

Standard Error of Est. = 6.45478 Mean absolute error = 5.33981

Durbin-Watson statistic = 0.0296894 (P=0.0000)




Figure 12k: Polynomial Regression Result for 1996 bathymetry

Multiple Regression Analysis for 1996 Bathymetry Survey Depth and Transformed River Coordinates Note: U = Cross Flow Coordinate; V = Along Flow Coordinate)

0.4037

0.6864

Dependent variable: Depth

 $\Delta * \Delta$

Parameter	Estimate	Standard Error	T Statistic	P-Value
CONSTANT	-16.90500	0.899662	-18.7904	0.0000
U	-0.36220	0.318288	-1.1380	0.2551
V	0.00854	0.008793	0.9716	0.3313
U*U	5.16838	0.095408	54.1717	0.0000

U*V -0.00683 0.001346 -5.0771 0.0000

0.000021

Analysis of Variance

0.00001

Source	Sum of Squares	Df	Mean Square	F-Ratio	P-Value
Model Residual	149683.0 747140.0	5 18373	29936.6 40.6651	736.17	0.0000

Total (Corr.) 896823.0 18378

R-squared = 16.6904 percent

R-squared (adjusted for d.f.) = 16.6677 percent

Standard Error of Est. = 6.37692

Mean absolute error = 5.26645

Durbin-Watson statistic = 1.95342 (P=0.0008)

Figure 12I: Polynomial Regression Result for 1997 bathymetry

Multiple Regression Analysis for 1997 Bathymetry Survey Depth and Transformed River Coordinates (Note: $U = Cross\ Flow\ Coordinate;\ V = Along\ Flow\ Coordinate)$

Dependent variable: Depth

Standard T

Parameter	Estimate	Standard Error	Statistic	P-Value
CONSTANT	-15.43800	0.889624	-17.3534	0.0000
υ	0.41462	0.327394	1.2664	0.2054
V	-0.00962	0.008684	-1.1077	0.2680
U*U	4.62971	0.098426	47.0377	0.0000
Λ*Δ	0.00006	0.000020	2.8574	0.0043
U* V	-0.00960	0.001377	-6.9715	0.0000

Analysis of Variance

Source	Sum of Squares	Df	Mean Square	F-Ratio	P-Value
Model Residual	122616.0 696994.0	5 17970	24523.2 38.7865	632.26	0.0000
m , 1 (a)	040640	48085			

Total (Corr.) 819610.0 17975

R-squared = 14.9603 percent

R-squared (adjusted for d.f.) = 14.9366 percent

Standard Error of Est. = 6.22788

Mean absolute error = 5.1599

Durbin-Watson statistic = 1.46572 (P=0.0000)

Figure 12m: Polynomial Regression Result for 1999 bathymetry

Multiple Regression Analysis for 1999 Bathymetry Survey Depth and Transformed River Coordinates (Note: U = Cross Flow Coordinate; V = Along Flow Coordinate)

Dependent variable: Depth

Parameter	Estimate	Standard Error	T Statistic	P-Value
CONSTANT	-23.76950	0.87765	-27.0832	0.0000
υ	-9.85349	0.31098	-31.6848	0.0000
V	0.08111	0.00884	9.1712	0.0000
U*U	7.05628	0.09114	77.4202	0.0000
V*V	-0.00019	0.00002	-8.9664	0.0000
Π*Δ	0.04073	0.00137	29.702	0.0000

Analysis of Variance

Source	Sum of Squares	Df	Mean Square	F-Ratio	P-Value
Model Residual	193813.0 471533.0	5 15 44 3	38762.6 30.5338	1269.50	0.0000

Total (Corr.) 665346.0 15448

R-squared = 29.1296 percent

R-squared (adjusted for d.f.) = 29.1067 percent

Standard Error of Est. = 5.52574

Mean absolute error = 4.52886

Durbin-Watson statistic = 1.59393 (P=0.0000)

Figure 12n: Polynomial Regression Result for 2001 bathymetry

Multiple Regression Analysis for 2001 Bathymetry Survey Depth and Transformed River Coordinates (Note: $U = Cross\ Flow\ Coordinate;\ V = Along\ Flow\ Coordinate)$

Dependent variable: Depth

		Standard	T	
Parameter	Estimate	Error	Statistic	P-Value
CONSTANT	-26.8546	0.86168	-31.1653	0.0000
U	-10.7103	0.30563	-35.0432	0.0000
V	0.1128	0.00868	12.9854	0.0000
บ*บ	6.7680	0.08866	76.3319	0.0000
V*V	-0.0003	0.00002	-12.4402	0.0000
U*V	0.0447	0.00134	33.2957	0.0000

Analysis of Variance

Source	Sum of Squares	Df	Mean Square	F-Ratio	P-Value
Model Residual	194485.0 468355.0	5 15805	38896.9 29.6334	1312.61	0.0000
Total (Corr.)	662840.0	15810			

R-squared = 29.3411 percent
R-squared (adjusted for d.f.) = 29.3188 percent
Standard Error of Est. = 5.44365

Mean absolute error = 4.47725

Durbin-Watson statistic = 1.59144 (P=0.0000)

Figure 12o: Polynomial Regression Result for 2002 bathymetry

Multiple Regression Analysis for 2002 Bathymetry Survey Depth and Transformed River Coordinates (Note: U = Cross Flow Coordinate; V = Along Flow Coordinate)

Dependent variable: Depth

Parameter	Estimate	Standard Error	T Statistic	P-Value
CONSTANT	-15.1029	0.481731	-31.3513	0.0000
ΰ	2.3383	0.233236	10.0253	0.0000
V	0.0291	0.005260	5.5312	0.0000
U*U	1.1821	0.094716	12.4807	0.0000
Λ*Δ	-0.0001	0.000013	-8.4951	0.0000
U*Δ	-0.0176	0.001022	-17.2185	0.0000

Analysis of Variance

Source	Sum of Squares	Df	Mean Square	F-Ratio	P-Value
Model Residual	65099.7 627880.0	5 18963	13019.9 33.1108	393.22	0.0000

Total (Corr.) 692980.0 18968

R-squared = 9.39416 percent

R-squared (adjusted for d.f.) = 9.37027 percent

Standard Error of Est. = 5.7542

Mean absolute error = 4.7017

Durbin-Watson statistic = 0.0507036 (P=0.0000)

Figure 12p: Polynomial Regression Result for 2004 bathymetry

Multiple Regression Analysis for 2004 Bathymetry Survey Depth and Transformed River Coordinates (Note: $U = Cross\ Flow\ Coordinate;\ V = Along\ Flow\ Coordinate)$

Dependent variable: Depth

Parameter	Estimate	Standard Error	T Statistic	P-Value
CONSTANT	-13.6587	0.164017	-83.2761	0.0000
U	-1.95215	0.136465	-14.3052	0.0000
V	0.03363	0.002168	15.5093	0.0000
U*U	4.23460	0.076119	55.6312	0.0000
Λ*Δ	-0.00016	0.000007	-24.7688	0.0000
U*V	-0.00185	0.000639	-2.8930	0.0038

Analysis of Variance

Source	Sum of Squares	Df	Mean Square	F-Ratio	P-Value
Model Residual	207735.0 937440.0	5 24889	41546.9 37.6648	1103.07	0.0000

Total (Corr.) 1.14517E6 24894

R-squared = 18.14 percent

R-squared (adjusted for d.f.) = 18.1236 percent

Standard Error of Est. = 6.13717

Mean absolute error = 5.03368

Durbin-Watson statistic = 0.0500437 (P=0.0000)

Figure 13: Estimated trend surface for 1995 bathymetry elevations. Note that the long- and cross-flow coordinates are not to scale or proportion.





Figure 14a: Directional Semivariograms and Fitted Models for Normal Scores Transformed Residuals of Detrended 1989 Bathymetry







Figure 14b: Directional Semivariograms and Fitted Models for Uniform Scores
Transformed Residuals of Detrended 1989 Bathymetry







Figure 14c: Directional Semivariograms and Fitted Models for Normal Scores Transformed Residuals of Detrended 1995 Bathymetry







Figure 14d: Directional Semivariograms and Fitted Models for Uniform Scores Transformed Residuals of Detrended 1995 Bathymetry







Figure 14e: Directional Semivariograms and Fitted Models for Normal Scores
Transformed Residuals of Detrended 1996 Bathymetry







Figure 14f: Directional Semivariograms and Fitted Models for Uniform Scores Transformed Residuals of Detrended 1996 Bathymetry







Figure 14g: Directional Semivariograms and Fitted Models for Normal Scores
Transformed Residuals of Detrended 1997 Bathymetry







Figure 14h: Directional Semivariograms and Fitted Models for Uniform Scores
Transformed Residuals of Detrended 1997 Bathymetry







Figure 14i: Directional Semivariograms and Fitted Models for Normal Scores Transformed Residuals of Detrended 1999 Bathymetry







Figure 14j: Directional Semivariograms and Fitted Models for Uniform Scores
Transformed Residuals of Detrended 1999 Bathymetry







Figure 14k: Directional Semivariograms and Fitted Models for Normal Scores Transformed Residuals of Detrended 2001 Bathymetry







Figure 14l: Directional Semivariograms and Fitted Models for Uniform Scores
Transformed Residuals of Detrended 2001 Bathymetry







Figure 14m: Directional Semivariograms and Fitted Models for Normal Scores Transformed Residuals of Detrended 2002 Bathymetry







Figure 14n: Directional Semivariograms and Fitted Models for Uniform Scores
Transformed Residuals of Detrended 2002 Bathymetry







Figure 14o: Directional Semivariograms and Fitted Models for Normal Scores Transformed Residuals of Detrended 2004 Bathymetry







Figure 14p: Directional Semivariograms and Fitted Models for Uniform Scores
Transformed Residuals of Detrended 2004 Bathymetry






Figure 15: Three realizations of the 1996 bathymetric elevations.






Figure 16: Histogram of sample elevations (Panel A) and simulated elevations (Panel B) for the lower Passaic River in 1995.



Note:

- 1) Simulated elevations represent all locations at which values were simulated, as opposed to the subset of locations at which inter year comparisons were conducted.
- 2) This comparison illustrates that the simulation algorithm reproduces the data histogram.
- 3) Because inter-year comparisons of simulated data were restricted to a smaller lateral extent than the sample data, direct comparison of histograms would be biased toward deeper soundings in the simulated soundings.

Figure 17: Semivariograms for 20 realizations compared with theoretical model semivariograms for cross-flow (green) and long-flow (black) directions.



a Minimum 70% Level of Confidence

P:\0285924\Mapping\CSM_MassBalance\bathymetry_frequency_depth_map_

a Minimum 70% Level of Confidence Lower Passaic River Restoration

a Minimum 70% Level of Confidence

P:\0285924\Mapping\CSM_MassBalance\bathymetry_frequency_depth_map_

a Minimum 70% Level of Confidence Lower Passaic River Restoration

P:\0285924\Mapping\CSM_MassBalance\bathy_simulation_deposition_v2.mxd

P:\0285924\Mapping\CSM_MassBalance\bathy_simulation_deposition_v2.mxd

P:\0285924\Mapping\CSM_MassBalance\bathy_simulation_deposition_v2.mx

at a Minimum 70% Level of Confidence in Non-Significant Erosion Areas Lower Passaic River Restoration

J:\1work\Passaic\bathy_simulation_deposition.mxd

at a Minimum 70% Level of Confidence in Non-Significant Erosion Areas Lower Passaic River Restoration

P:\0285924\Mapping\CSM_MassBalance\bathy_simulation_deposition_v2.mxd

Elevation Changes in Non-Significant Erosion Areas Lower Passaic River Restoration

P:\0285924\Mapping\CSM_MassBalance\bathy_simulation_deposition_v2.mxc

Elevation Changes in Non-Significant Erosion Areas Lower Passaic River Restoration

Elevation Changes in Non-Significant Erosion Areas Lower Passaic River Restoration

P:\0285924\Mapping\CSM_MassBalance\bathy_simulation_deposition_v2.mx

Elevation Changes in Non-Significant Erosion Areas

Lower Passaic River Restoration




Figure 23b Histogram of 2007 Multibeam - 2007 Single Beam Cross Transects









Figure 23c Histogram of 2007 Multibeam - 2007 Single Beam Longitudinal Transects

























































Figure 26a: Conditional Simulation Validation:
Difference between 2007 Actual Multibeam Surface
and Predicted 2007 Surface




Figure 26b: Passaic Bathymetric Change Observations






Figure : Conditional Simulation Validation False Positive rates

2007 Multibeam Surface Approximated by the 1995 Locations via Conditional Simulation

Threshold	P(Erode>12-in)	P(Erode>6 in)	P(Erode>3 in)	P(Erode>0 in)
>31% probability	5%			
>50% probability	2.09%	5.23%	12%	49.7%
>62.5% Probability			5%	
>70% probability	1.178%	2.156%	3.367%	6.556%
>74% probability				5%












Figure 29: Segments of the Passaic River where 2008 and 2007 multibeam bathymetry data were compared based on average depth in a 3 ft by 3 ft grid spacing.













Figure 30c:

River Segment A (RM13.6) Transect bathymetry in 2007 and 2008 and change (2008-2007).

Figure 30d: Difference between 2007 and 2008 multibeam bathymetry (feet; negative deeper in 2008) for River Segment A (RM13.6) with transect location



sand waves were eroded between 2007 and 2008 surveys N S





Figure 31c: River Segment B (RM9.75) Transect bathymetry in 2007 and 2008 and change (2008-2007).

Depths (x-axis) and Distances (y-axis) in Feet

Coarse bed beneath bridge is about 0.2 feet deeper in 2008 than in 2007. It is likely that there was no erosion beneath bridge.








Figure 32c: River Segment C (RM8.85) Transect bathymetry in 2007 and 2008 and change (2008-2007).





NW SE





Figure 33c: River Segment D (RM4.05) Transect bathymetry in 2007 and 2008 and change (2008-2007).







Figure 34c: River Segment E (RM1.6) Transect bathymetry in 2007 and 2008 and change (2008-2007).






Figure 35. Net elevation change from 1989 through 2007.